

- Growth of semi-permeable membrane of defined geometry  
- by **flowing gas** and **liq.** nutritional phases over  
substrate having cell layer which grows in  
controlled manner and is removed for further use,  
e.g. in human organ transplantation stations.

L20 ANSWER 92 OF 121 WPINDEX COPYRIGHT 2002 DERWENT INFORMATION LTD  
AN 1995-240685 [31] WPINDEX  
DNC C1995-110405  
DC D16 D22  
IN PRENOSIL, J E  
PA (PREN-I) PRENOSIL J E  
CYC 2  
PI WO 9517526 A1 19950629 (199531)\* DE 22p  
AU 9511900 A 19950710 (199543)  
CH 687153 A5 19960930 (199644)  
ADT WO 9517526 A1 WO 1994-CH241 19941222; AU 9511900 A AU 1995-11900 19941222;  
CH 687153 A5 CH 1993-3830 19931222  
FDT AU 9511900 A Based on WO 9517526  
PRAI CH 1993-3830 19931222  
AN 1995-240685 [31] WPINDEX  
AB WO 9517526 A UPAB: 19950810

The growth of a semi-permeable membrane uses a reaction chamber (1) and a medium chamber (2) through which a gas phase and a liquid nutritional phase flow. They pass over a semi-permeable membrane (M1) which serves as the substrate for the new membrane. The novelty is that the new cell material is inoculated to the semi-permeable membrane to form a first layer. The lower side of the semi-permeable membrane is flushed by nutritional solution and the upper side is flushed by the gas phase. Controlled cell growth ensues to form a number of layers and a cell structure (ML) from which metabolic by-products are removed by the gas and liquid phases to produce a cell structure which can be removed under sterile conditions for further use.

USE- The process and assembly produce a semi-permeable membrane of defined geometry, as used in the stations associated with human organ transplantation.

ADVANTAGE - The process and assembly are suitable for automated operation, thus largely removing the requirement for manual intervention.  
Dwg.1/3